## **CUMULATIVE INDEX 1997**

## Volume 14

PEDIATRIC PODIATRY, pages 1-233 January

April IMAGING OF THE FOOT AND ANKLE, pages 235–405

July SPORTS MEDICINE AND REHABILITATION, pages 407–584 October AMPUTATIONS OF THE LOWER EXTREMITY, pages 585–820

Note: Page numbers of article titles are in boldface type.

Abscess, in diabetes mellitus, 240-241 magnetic resonance imaging of, 254-256 treatment of, 673-674, 676-677 Acetabular dysplasia. See Hip, instability of.

Acetabular index, in hip radiology,

190-191

Achilles tendon, imaging of, in inflammation, 289-292

in rupture, 317, 321

magnetic resonance, 317-320

lengthening of, in flatfoot, 96-97 with midfoot amputation, 756-759

Achilles tendon injuries, in athletes, 447-458

classification of, 448-449

diagnosis of, 449-452 disability in, 448

etiology of, 447-448

incidence of, 447

insertional tendonopathy, 449-450

midportion, 451-452 myotendinous junction, 452

pathology of, 448

prevention of, 449

treatment of, 450-453

young, 574

in ballet dancers, 528-529

Acupuncture, in peroneal tendon injuries,

Adjuvant therapy, with amputation, in

malignancy, 615-616 Airway obstruction, postoperative, in

pediatric patients, 146 Albumin levels, in healing prediction, 670,

Ambulation, after amputation, metabolic cost of, 599-602

midfoot, 759-760

pathomechanics of, 776-777 potential for, 594-596

transmetatarsal, 759-760

weight distribution in, 731-732

biomechanics of, 775-776 Americans with Disabilities Act,

amputation and, 585-586

Amniotic band sequence (constricting

band syndrome), 633-636 Amputation, 585-597

Achilles tendon lengthening with, 756-

ambulation after. See Ambulation.

Americans with Disabilities Act and, 585-586

art of, 667-678

at tarsometatarsal joints (Lisfranc), 753-

755, 763, 780-781 bilateral, metabolic cost of, 601

biomechanical effects of, 777-778

body image in, 587

Boyd-Pyrogoff, 763-764

Chopart, 754, 756, 763

prostheses for, 780-782

complications of, 590

concurrent health problems with, 594

contralateral foot condition in, 593

discharge plan for, 595

emotional shock in, 589-590

failure of, 728-729

hallux, 720, 777, 779

health care team for, 586

in infection or ischemia, 667-689

in lawnmower injury, 605-608

in pediatric patients. See Pediatric pa-

internal versus external, 682, 684 level of, 587-588

Amputation (Continued) in infection or ischemia, 668-671, 681-682 midfoot, 742-744 transmetatarsal, 742-744 Lisfranc and modifications, 753-755, 763 prostheses for, 780-781 long-term care after, 785-800 metabolic cost of, 599-602 midfoot, 741-762 phantom sensation and pain in, 589, 591 postoperative care in. See Postoperative preoperative preparation for, 588 preventive interventions for, 785-800 Firm system in, 790-792 Preservation Amputation Care and Treatment program in, 787-790 primary care provider in, 786 priorities for, 786 smoking cessation programs in, 792toes, 692-693 prostheses for. See Prostheses. ray resections and. See Ray(s), resection recreational activities in, 596 rehabilitation in. See Rehabilitation. repeat, prevention of. See Amputation, preventive interventions for. stump care in, 592-593 support groups for, 597 Syme's. See Syme's amputation. toe. See under Toe(s). transmalleolar. See Syme's amputation, transmalleolar. transmetatarsal. See Transmetatarsal amputation. tumor as indication for, 613-620 using freezing, 676-677 versus reconstruction, in major trauma, Analgesia, in pediatric patients, 131-148 complications of, 143-144 definition of, 132 nothing by mouth guidelines for, 135 opioids in, 136 outside operating room, 132-133 postoperative, 144-147 preoperative examination for, 133-134 Anaphylactic reactions, perioperative, 144 Anchoring systems, for arthroscopic ankle stabilization, in athletes, 471-473,

475-476

Anesthesia, for amputation, 745

in pediatric patients, 131-148

complications of, 143-144

controversies in, 138-139

definition of, 131-132

Syme's, 765-766

epidural, 141 fluid management in, 140 intraoperative, 139-143 intravenous lines for, 140 nothing by mouth guidelines for, 135 outpatient, 134-135 outside operating room, 132-133 postoperative considerations in, 144-147 premedications and, 136-138 preoperative examination for, 133-134 regional, 141 selection of, 139-140 spinal, 141 Aneurysmal disease, angiography of, 270-274 Angiography, of adventitial cystic disease, 273 of arterial aneurysmal disease, 270-274 of arterial entrapment, 273 of arterial occlusive disease, 268-270 of diabetic foot, 244-245 of ergotism, 273 of thromboangiitis obliterans, 273 of venous disease, 278-279 Ankle, amputation and salvage procedures in, in congenital conditions, 621-633 fibular hemimelia, 623-624, 626-632 tibial hemimelia, 624-627, 632-633 arthroscopic stabilization of, in athletes, 459-478 advantages of, 468 anatomic considerations in, 460-461 anchoring systems for, 471-473, 475-476 diagnosis before, 462, 470 pathomechanics of, 460-461 portals for, 469 techniques for, 468-474 versus nonarthroscopic procedures, 462-468 deformity of, in clubfoot, 18, 20 disarticulation of, in tibial hemimelia, Syme's. See Syme's amputation, transmalleolar injuries of, in ballet dancers, 529 in young athletes, 570-573 sprains of, arthroscopic stabilization of. See Ankle, arthroscopic stabilization of. rehabilitation in, 539-555 Ankle block anesthesia, for amputation, Ankle-brachial index, in healing prediction, 668-669, 765 Ankle-foot orthosis, for midfoot amputation, 781 Ankylosing spondylitis, imaging of, 374 Ankylosis, in arthritis, radiography of, 371 Anterior drawer test, in ankle instability,

Anteroposterior view, in radiography, of calcaneal fractures, 341
Antetorsion, definition of, 58
femoral, in-toe gait in, 61–68
Anteversion, hip, normal angle of, 63–64
Anteversion syndrome, 58, 60
Antibiotics, in amputation, 699, 726
in osteomyelitis, 675–676
preoperative, 135–136
Antibodies, technetium 99m-labeled, in

diabetic foot scan, 250, 252 Anti-inflammatory agents, homeopathic, in Achilles tendon injuries, 450-451,

in peroneal tendon injuries, 456–457 Antiperspirants, in hyperhidrosis, in athletes, 515–516

Aorta, abdominal, angiography of, in aneurysmal disease, 270–272 in occlusive disease, 268–269

Aortography, of aneurysmal disease, 270–272

Apophyseal injures, in young athletes, 573–574

Arterial calcification, in diabetes mellitus, 237, 240

Arterial occlusion, amputation and salvage procedures in, 659

Arthritis, gouty, heel pain in, 295-296 imaging of, 379-381

imaging of, **387–398**. See also Arthritis, radiography of.

arthrography in, 389-391 bone scanning in, 387-389

computed tomography in, 392, 394-396

magnetic resonance, 395, 397 tenography in, 390, 392–393 radiography of, 369–385

degenerative (osteoarthritis), 377–379 gouty (crystal deposition), 379–381 in connective 1882 diseases, 381–382

neuropathic, 382–384 osteomyelitis, 382–384 principles of, 369–370

psoriatic, 374-376 rheumatoid, 372-374

septic, 382–384

seronegative spondyloarthropathies, 374–377

sesamoid bones, 384 rheumatoid, bone scan of, 389

heel pain in, 295 imaging of, 372–374

septic, imaging of, 382–384 Arthrodesis, of first metatarsophalangeal joint, in hallux limitus, 425

Arthrography, of arthritis, 389–391 Arthropathy, Charcot, after amputation, 731, 733–734 Arthroplasty, as amputation alternative, 694–695, 702–707 implant, in hallux limitus, 425 Arthrorisis, subtalar, 87–98

Arthroscopy, in ankle stabilization. See Ankle, arthroscopic stabilization of.

Athletes. See Sports medicine.

Athlete's foot, 517-519

Atrophic neuropathy, in diabetes mellitus, 236-237

Austin procedure, in hallux valgus, 438–441

Austin-Youngswick procedure, in hallux limitus, 424

Automated profile testing, preoperative, in pediatric patients, 168–172 Avascular necrosis, of calcaneus, imaging

of, 288–289

Bacterial culture, of amputation site, 697 Balance, with amputation, 594 Ballet, overuse injuries in, **525–532** Barlow's test, for hip instability, 188 Baths, contrast, in athlete rehabilitation, 535

BeBax shoe, 210 Beighton-Horan criteria, for ligamentous laxity, 125, 127

Biomechanics, of ambulation, 775–776 after amputation, 777–778

Biopsy, of tumor, 615 Black heel syndrome, in athletes, 511–512 Blisters, in athletes, 512

Blood testing, preoperative, in pediatric patients, 164–171

Blood vessels, transection of, in amputation, 697

Body image, in amputation, 587 Bohler's angle, 339

Bone, debridement of, as amputation alternative, 701

sclerosis of, in arthritis, radiography of, 371

Bone graft, in lawnmower injury reconstruction, 606

Bone scans. See Nuclear medicine scans. Bonney-Kessel procedure, in hallux limitus, 424–425

Boots, compression, in athlete rehabilitation, 538 cryoamputation, 676–677

Boyd-Pyrogoff amputation, 763–764 Brachman skate, 225 Brachymetatarsia, 642–643

Bracing. See Splinting and bracing. Broden's view, in radiography, of calcaneal fractures, 341

Brostrom procedure, for ankle stabilization, 467

Brown's technique, in tibial hemimelia treatment, 632–633
Bruises, in athletes, 512–513
Buerger's disease, angiography of, 273
Bunion. See *Hallux valgus*.
Bupivacaine, in pediatric surgery, 140–141
Bursal cysts, in athletes, 523
Bursitis, Achilles, imaging of, 289–292, 319
in athletes, 513
retrocalcaneal, imaging of, 291–292, 319–320

Calcaneal apophysis, osteochondrosis of (Sever's disease), 298 Calcaneal petechiae (black heel syndrome), in athletes, 511-512 Calcaneocuboid joint, deformity of, in clubfoot, 20, 23 Calcaneofibular ligament, anatomy of, 460 pathomechanics of, 461 reconstruction of, 463-467 repair of, arthroscopic, 468-474 surgical, 467 Calcaneovalgus deformity, in hip instability, 183 shoe therapy for, 212 Calcaneus, avascular necrosis of, imaging of, 288-289 changes in, in Reiter's syndrome, 377 in rheumatoid arthritis, 373-374 deformity of, in clubfoot, 18 fractures of, 337-356 anatomic considerations in, 338-339 anterior process, radiography of, 342 avulsion, radiography of, 342 beak, radiography of, 342 classification of, 337 imaging of, 287-288 computed tomography in, 344-353 magnetic resonance, 353 radiography in, 341-343 radionuclide scintigraphy in, 353 mechanism of injury in, 340 old, computed tomography of, 351 stress, 287-288 radionuclide scans of, 353 tuberosity, radiography of, 342 pain within. See Heel-spur syndrome. tumors of, imaging of, 292-293

Calcification, arterial, in diabetes mellitus, 237, 240
Calcium pyrophosphate dihydrate crystal deposition disease, imaging of, 379
Cancer. See Malignancy.
Candidiasis, of nail, in athletes, 517
Cast, in clubfoot treatment, 4–6, 32–35

postoperative, for amputation, 588-589 Cavus foot, in Charcot-Marie-Tooth disease, 103 in hereditary motor sensory neuropathy, pathogenesis of, 112
C-E angle, in hip radiology, 192
Cellulitis, in diabetes mellitus, 240–241 magnetic resonance imaging of, 254–256 nuclear medicine scans of, 247
Central polydactyly, 655–657
Cerebral palsy, hip instability in, 184–185, 197
Charcot arthropathy, after amputation, 731, 733–734

Charcot arthropathy, after amputation, 731, 733–734
Charcot-Marie-Tooth disease, 99–116 electrodiagnostic studies of, 110–111 historical review of, 100 in pediatric patients, 111 pathogenesis of, 111–112 treatment of, 112–113 type I (hypertrophic), 101–104 pathogenesis of, 109–110 type IA, 104–105 type IB, 105–106

type IA, 104–105
type IB, 105–106
type IC, 106
type II (neuronal form), 106–108
pathogenesis of, 109–110
subtypes of, 107
types X1 and X2, 106
Cheilectomy, in hallux limitus, 423–424
Chemotherapy, with amputation, in

malignancy, with amputation, in malignancy, 615-616
Chopart amputation, 754, 756
prostheses for, 780-782
versus Syme's ankle disarticulation, 763
Chopart joint, neuroarthropathy of, in

diabetes mellitus, 237, 239 Chrisman-Snook modification, of Elmslie procedure, for ankle stabilization, 463,

Chromosomal abnormalities, hip instability in, 184

Cigarette smoking cessation programs, in amputation prevention, 792–798
Circular Torquheel device, 227–228

Claudication, in peripheral vascular disease, abdominal aortic, 268–269 exercise program for, 267 femoral arterial, 269 iliac arterial, 269

Clawtoes, in hereditary motor sensory neuropathy, pathogenesis of, 112 Cleft foot deformity (ectrodactyly),

637-642 Closure, of amputation site, 699-700 Clubfoot, 9-56 causes of, 15 classification of, 14-16 clinical evaluation of, 23, 26

constricting bands with, 636 historical review of, 11–14 incidence of, 14 late presentation of, treatment of, 4-5 pathologic anatomy of, 15, 17-23 articular, 18, 20-23

osseous, 18 soft tissue, 18-19

radiographic evaluation of, 25, 27-30

treatment of, 31-43

casting in, 4-6, 32-35

Denis Browne splint in, 1-4 gangrene after, 659-660

historical review of, 1-7

in late presentation, 4-5 incisions for, 37

long-term results of, 43-52

shoes in, 209-211

soft tissue release in, 37-43 surgical, 35-43

timing of, 31-32, 36-37

Coagulation disorders, arterial occlusion in, 659

Coalition, rearfoot, 624

Cognitive deficits, in amputation, 595 Cold therapy, in athlete rehabilitation, 534-535

Colville procedure, for ankle stabilization,

Compartment syndrome, in athletes, 454, 575

in major trauma, 611

Complete blood count, in preoperative testing, in pediatric patients, 164-168

Compression, intermittent, in athlete rehabilitation, 538

Compression syndromes, venous dysfunction in, 278-279

Computed tomography, of arthritis, 392, 394-396

of calcaneal fractures, 344-353 classification system for, 350-353 technique for, 344-349

three-dimensional, 349-350

of diabetic foot, 244

of hip, 193

of Morton's neuroma, 305

of os trigonum syndrome, 297

of tarsometatarsal joint complex injury, 367

of tibial torsion, 71-72

Congenital conditions, amputation in. See Pediatric patients, amputation and salvage procedures in, in congenital conditions.

convex pes valgus, in hip instability, 183-184

shoe therapy for, 212

Connective tissue diseases, arthritis in, imaging of, 381-382

Constricting band syndrome, 633-638 Contact dermatitis, in athletes, 519 Contractures, in clubfoot, 18-19

Contralateral foot, in amputation, 593-595 Contrast baths, in athlete rehabilitation, 535

Convex pes valgus, congenital, in hip instability, 183-184

shoe therapy for, 212

Corns, in athletes, 522

Corynebacterium minutissimum infections, in athletes, 523

Cotton osteotomy, of cuneiform, in flatfoot, 96

Counter Rotation System, for in-toe gait correction, 81

Crosby-Fitzgibbons classification, of calcaneal fractures, 350

Crush injury, reconstruction in, 604-605 Cryoamputation, 676-677 Cryotherapy, in athlete rehabilitation,

534-535 Crystal deposition diseases, imaging of,

379-381 Culture, of amputation site, 697

Cuneiform, osteotomy of, in flatfoot, 96 Cysts, subchondral, in arthritis, radiography of, 371 synovial, in athletes, 523

Dance, overuse injuries in, 525-532

Dantrolene, in malignant hyperthermia, 143-144

Dead spaces, at amputation site, 697-699 Débridement, as amputation alternative,

in amputation, 700-701 in crush injury, 604-605

in infection, 670, 674-677, 680-681, 694 in lawnmower injury, 605

in trauma, 718

Deformities, congenital, treatment of. See Pediatric patients, amputation and salvage procedures in.

Degenerative joint disease, imaging of, 377-379

Degloving injuries, treatment of, 717-718 Dejerine-Sottas syndrome, 108

Denis Browne splints, 1-4, 223-224 in in-toe gait, 77-81

Dermatitis, in athletes, 519

Dermatofibromas, in athletes, 524 Dermatologic problems, in athletes,

511-524

black heel syndrome, 511-512 blisters, 512

bruises, 512-513

bursitis, 513

erythrasma, 523

excrescences, 521-523 fissures, 513-514

Dermatologic problems (Continued) foreign bodies, 514–515 ganglions, 523–524 hyperhidrosis, 515–516 nail diseases, 516–517 psoriasis, 519–521 puncture wounds, 514–515 tinea pedis, 517–519 in ballet dancers, 531

Dermatophytosis, in athletes, nail, 516–517 skin, 517–519

Desflurane, for pediatric patients, 142 Desyndactyly, 655

Dexamethasone, in iontophoresis, in athlete rehabilitation, 536

Diabetic foot, 235–264 contributing factors in, 235–241 imaging of, 241–261

after surgical changes, 259 angiography in, 244 computed tomography in, 244 in edema, 253–254

in ischemia, 258–259 in neuroarthropathy, 257–258, 261,

383–384 in osteomyelitis, 244, 246–252, 256– 257, 260, 383–384

in soft-tissue fluid collections, 254–256 magnetic resonance, 252–261 magnification radiography, 243–244

nuclear medicine scans in, 244, 246–252

radiography in, 242–243 in athletes, precautions in, 520–521 infection in, 240–241

ischemia and infection in. See Ischemia, amputation and salvage procedures in.

amputation and salvage procedures in.

long-term care of. See Amputation, preventive interventions for.

neuroarthropathy in, 235–239, 257–258,

261, 383–384

toe amputation in, 691–693 vascular compromise in, 237, 240 Digit(s). See *Toe*(s).

Digital nerves, neuroma of. See Morton's neuroma.

Disarticulation, ankle, in tibial hemimelia, 632

Syme's. See Syme's amputation, transmalleolar.

metatarsophalangeal joint, 715–717 Dislocation, hip. See Hip, instability of. Disseminated intravascular coagulation, ischemia in, amputation and salvage procedures in, 657

Doppler studies, in healing prediction, 669–670

Dorsal-plantar flaps, in digital amputation, 709–711

Drains, for amputation site, 697-698, 706

Dressings, for amputation, 588–589, 726–727 transmetatarsal, 747 for infected wounds, 670–671

Droperidol, as premedication, for pediatric patients, 137

Drugs, in athlete rehabilitation, in iontophoresis, 536 in phonophoresis, 537–538

Dry gangrene, treatment of, 678–681 Duplication, of toes (polydactyly), 652, 654–657

Dyskinesia, post-static, in heel-spur syndrome, 503

Dysplasia, hip. See Hip, instability of.

Eastwood classification, of calcaneal fractures, 350

Ecchymosis, in athletes, 512–513 Ectrodactyly, 637–642

Edema, in diabetes mellitus, magnetic resonance imaging of, 253–254 intermittent compression in, 538

EDIN procedure, in Morton's neuroma, in athletes, 497-500

Ehlers-Danlos syndrome, ligamentous laxity in, 125–126

Electrical agents, in athlete rehabilitation, 536–537

Electrodiagnostic studies, in hereditary motor sensory neuropathy, 110–111 type I, 103–104 type II, 107–108 type III, 108

Elmslie procedure, for ankle stabilization, 463, 466

Emotional shock, in amputation, 589–590 Energy cost, of amputation, 599–602 Enflurane, for pediatric patients, 142 Environmental modifications, for

amputation, 596 Epiphyseal arrest, plantar, of first metatarsal, in flatfoot, 96 Epiphysiodesis, in macrodactyly, 650

Epiphysiodesis, in macrodactyry, 60 Equinovalgus deformity, in fibular hemimelia, 624 in tibial hemimelia, 625

Equinovarus deformity, in midfoot amputation, 753–754

Achilles tendon lengthening in, 756–759

Ergotism, angiography of, 273 Erosions, in arthritis, radiography of, 371, 380–381

Erythrasma, in athletes, 523 Eschar removal, in amputation, 700 Evans procedure, for ankle stabilization, 463-464 Excrescences, in athletes, 521–523
Exercise, heel-spur syndrome development in, 508–509
in athlete rehabilitation, alternative programs for, 533–534
in ankle sprain, 539–555
in heel-spur syndrome, 556
in plantar fasciitis, 556
in sesamoiditis, 556
in turf toe, 555–556
return-to-activity, 543, 551, 554

in foot deformities, in hereditary motor sensory neuropathy, 112 in peripheral vascular disease, 267

Exostosis, retrocalcaneal, in athletes, 449–450

Extensor tendinitis, in ballet dancers, 527–528

Facial dysmorphisms, hip instability with, 184

Fascia lata graft, for ankle stabilization, 463, 466 Fasciitis, plantar, in athletes, rehabilitation

in, 556

young, 569 Femoral antetorsion, in-toe gait in, 61–68 residual, consequences of, 75–76

Femoral anteversion, definition of, 58–59 in-toe gait in, gait compensations in, 73–75

Femoral arteries, angiography of, in aneurysmal disease, 271, 273 in occlusive disease, 269

in occlusive disease, 269
Femoral torsion, in-toe gait in, 61–68
Denis-Browne splints for, 80–81
shoe therapy for, 211

splinting for, 218 Femur, displacement of, in hip radiology, 192

Fentanyl, in pediatric patients, 138 Fetus, hip instability in, 182 ligamentous laxity in, 118, 120

Fiberglass splints, 227 Fibromas, in athletes, 523–524

of hindfoot, imaging of, 293–294 Fibrosarcoma, treatment of, 616

Fibula, fractures of, in young athletes, 572 stress injuries of, in ballet dancers, 531

Fibular hemimelia, 623-624 treatment of, 626-632

Fillauer splint, 224

Firm system, in amputation prevention, 790–792

First ray, function of, in hallux limitus, 410 Fissures, in athletes, 513-514

Flatfoot, flexible, causes of, 124–125 clinical features of, 125, 127–128

in adolescents, 123-124

in infants, 120-121

in school-age children, 122-123

in toddlers, 121-122

intrauterine environment and, 118, 120

treatment of, 128

shoe therapy in, 212

subtalar arthrorisis in, 87-98

heel-spur syndrome with, 506 Flexible flatfoot. See Flatfoot, flexible.

Flexo splint, 225

Flexor hallucis longus tendon, injuries of, magnetic resonance imaging of,

328-329 tendinitis of, in ballet dancers, 526

Fold asymmetry, in hip instability, 189 Foot Risk Score, for diabetic foot, 787–788 Forefoot adductus, hallux limitus in, 410

Foreign bodies, in athletes, 514-515 Formaldehyde, in hyperhidrosis, in

athletes, 516

Fractures, after amputation, 731 calcaneal. See Calcaneus, fractures of.

in young athletes, 561

ankle, 572-573

fibular, 572

Jones', 570

malleolus, 572

tibial, 574 open, treatment of, 717-719

stress. See Stress fractures. Freezing, in amputation, 676–677

Freiberg's disease, in young athletes, 566 Friction blisters, in athletes, 512

Friedman counter splint, 225

Fungal infections, in athletes, nail, 516-517 skin, 517-519

Gait. See also Ambulation.

in-toe, 57-85

assessment of, 76-77

causes of, 60-77

hip rotation, 61–68 tibial torsion, 68–73

definitions of, 58-60

ternitions or, 58-60

transverse plane deformities in, 73-77 treatment of, 77-82, 209-212

out-toe, shoe therapy for, 212-213

Gait plates, for in-toe gait correction, 81-82 Gallium 67 scans, of diabetic foot, 247, 249

Ganglions, in athletes, 523–524

in heel, imaging of, 293 Gangrene, after clubfoot surgery, 659-660

dry, treatment of, 678-681 in arterial occlusion, 659

in purpura fulminans, 657-658

transmetatarsal amputation in, 742 Ganley splint, 224-225 Genu valgum, in-toe gait in, 211–212
Genu varum, in-toe gait in, 212
Giant cell tumors, of calcaneus, imaging of, 293
Gigantism, of toes, 644–652
Gissane's angle, 339
Glycosamine sulfate, in Achilles tendon injuries, 450, 453
Gouty arthritis, heel pain in, 295–296 imaging of, 379–381
Granulomas, foreign body, in athletes, 514–515
Gruca's procedure, in fibular hemimelia, 632
Guillotine (open) transmetatarsal amputation, 752–753

Haglund's disease, imaging of, 291-292 Hallux, abducto-valgus deformity of, after amputation, 731 amputation of, biomechanical effects of, prosthesis for, 779 transplantation of, to hand, 720 Hallux limitus, in athletes, 407-426 anatomic considerations in, 408 biomechanics of, 408-414 causes of, 408 classification of, 414-415, 418-419 clinical features of, 415-417 differential diagnosis of, 421-422 first-ray function in, 410 high gear-low gear theory of, 411 historical review of, 407-408 joint motion measurement in, 415-416 metatarsal shape and length in, 410-411 radiography of, 418-420 soft-tissue causes of, 412 treatment of, 422-425 Hallux rigidus, imaging of, 378 in young athletes, 566 Hallux set angle measurement, in hallux valgus, in athletes, 437-438 Hallux valgus, in athletes, correction of,

427–445

biomechanical evaluation for, 429–431 expectations in, 428

hallux set angle measurement in, 437–438 imaging for, 431–438

joint shape and condition evaluation before, 436–437 osteotomies in, 438–443 proximal articular set angle measurement in, 437 rehabilitation after, 443–445 Halothane, for pediatric patients, 142
Hamartomas, macrodactyly with, 645
Hammertoes, in hereditary motor sensory
neuropathy, pathogenesis of, 112
Hands-on-a-clock method, in hip range of
motion measurement, 66
Harpoon Anchor System, for arthroscopic

young, 566, 568

Harpoon Anchor System, for arthroscopic ankle stabilization, 476

Healing, amputation level selection and, 668–671 midfoot, 743–744

Syme's ankle disarticulation, 765 in pediatric patients, 561 Heat therapy, in athlete rehabilitation, 535

Heel, biomechanics of, 283–284 fissures of, in athletes, 513–514 Heel pad, migration of, after Syme's amputation, 769

Heel pain, 281–301 anatomic considerations in, 281–284 in Achilles tendinitis and bursitis, 289–

in Achilles tendinitis and bursitis, 289 292 in avascular necrosis, 288–289 in inflammation, 294–296

in os peroneum syndrome, 296–297 in os trigonum syndrome, 297–298 in Paget's disease, 297

in plantar fasciitis, 284–287 in reflex sympathetic dystrophy, 298–299 in Sever's disease, 298

in trauma, 287-288 tumor-related, 292-294

Heel-spur syndrome, in athletes, **503–509** alternative medicines in, 509 anatomic considerations in, 503–504 biomechanics of, 504–505 clinical examination in, 506–507 etiology of, 504–505 in pes cavus, 506 in therapeutic exercise programs,

508-509 radiography in, 507-508 rehabilitation in, 556 treatment of, 505-506

occupational, 508 Heloma, in athletes, 522

Hemihypertrophy, congenital, 644–652 of toes (macrodactyly), 644–652

Hemimelia, fibular, 623–624, 626–632 tibial, 624–627, 632–633 Hemorrhage, subcutaneous (black heel

syndrome), in athletes, 511–512 Hereditary motor sensory neuropathy,

electrodiagnostic studies of, 110–111 historical review of, 100 pathogenesis of, 109–112 treatment of, 112–113 type I (hypertrophic), 101–104 pathogenesis of, 109–110
type IA, 104–105
type IB, 105–106
type IC, 106
type II (neuronal form), 106–108
pathogenesis of, 109–110
subtypes of, 107
type III (Dejerine-Sottas syndrome), 108
type IV (Refsum's syndrome), 108–109
type V (spastic paraplegia), 109
type VI, 109
type VII (retinitis pigmentosa), 109
types X1 and X2, 106
Hereditary neuropathy with liability for
pressure palsy, 105

anteversion of, normal angle of, 63–64 femoral torsion at, in-toe gait in, 61–68 instability of, in pediatric-patients, 179–

Hilgenreiner's line, in hip radiology, 190 Hip, abduction limitation in, in instability,

208
case studies of, 197–206
clinical features of, 186–189, 194–197
congenital nature of, 180
imaging of, 190–193
incidence of, 182
natural history of, 193–194
pathologic anatomy of, 180–181
risk factors for, 182–186
screening for, 186
types of, 181

ligamentous laxity at, in in-toe gait, 64 Home equipment, in amputation rehabilitation, 593–594

Homeopathic remedies, in Achilles tendon injuries, 450–451, 453 in heel-spur syndrome, 509

in peroneal tendon injuries, 456–457 Hyperhidrosis, in athletes, 515–516

Hyperkeratosis, in athletes, 522 fissures in, 513-514 in hallux limitus, 416 plantar, 522-523

Hypermobile flatfoot, treatment of, subtalar arthrorisis in, 87–98

Hyperthermia, malignant, in pediatric patients, 143–144
Hypertrophic neuropathy, in diabetes

mellitus, 236–238 of infancy (Dejerine-Sottas syndrome),

Hypothermia, perioperative, in pediatric patients, 143

Ice therapy, in athlete rehabilitation, 534–535 Iliac arteries, angiography of, in aneurysmal disease, 271 in occlusive disease, 269 Imaging. See also specific modalities, e.g., Magnetic resonance imaging.

of arthritis, 369–385 special techniques for, 387–398 of calcaneal fractures, 337–356

of calcaneal fractures, 337–356 of diabetic foot, 235–264 of heel pain, 281–301

of Morton's neuroma, 303-311

of peripheral vascular disease, 265-280 of plantar faccilitis 281, 301

of plantar fasciitis, 281-301 of tarsometatarsal joint con

of tarsometatarsal joint complex injuries, 357–368

of tendon injuries, 313-335

Immunocompetence, in healing prediction, 670

Immunodeficiency, infection response in, 675–676

Immunoglobulins, indium 111-labeled, in diabetic foot scan, 250 Implant arthroplasty, in hallux limitus, 425 Incisions, for amputation, 696–697. See

also specific procedure. Indium 111 scans, of diabetic foot, 249–250 Infections, after amputation, management

of, 700–701 Syme's, 768

amputation and salvage procedures in, 667–689

art of, 667–668
control during procedure, 696
failure of, 728–729
internal versus external, 682–684
level for, 668–671, 681–682
postoperative care in, 684–686
procedure for, 672–677
team approach to, 671–672
transmetatarsal, 742
wound healing considerations in, 668–

bacterial, in tinea pedis, 518 débridement in, 694 in diabetes mellitus, 240–241 computed tomography of, 244 purpura fulminans in, amputation and salvage procedures in, 657–658 Inflammation, heel pain in, 294–296 synovial, imaging of, 372–377 Infracalcaneal heel pain. See Heel-spur

syndrome.

Injection therapy, for Morton's neuroma,

494–495 Injuries. See also *Trauma*.

in sports medicine. See Sports medicine. Interdigital neuroma. See Morton's neuroma.

Interferential stimulation, in athlete rehabilitation, 536–537

Intermetatarsal ligaments, transverse, interdigital nerve entrapment at. See Morton's neuroma. Intermetatarsal neuroma. See Morton's neuroma.

Intermittent compression, in athlete rehabilitation, 538

Interphalangeal joints, arthroplasty of, as amputation alternative, 702-705 first, dysfunction of, in hallux limitus, 416

Intertarsal joints, neuroarthropathy of, in diabetes mellitus, 236

In-toe gait, 57-85 causes of, 60-77 hip rotation, 61-68 tibial torsion, 68-73 definitions of, 58-60

hip motion measurement in, 65-68 radiography in, 66-68 spontaneous resolution of, 57-58 transverse plane deformities in, gait as-

sessment in, 76-77 gait compensations in, 73-75 residual, 75-76

treatment of, 77-82, 209-212 Iontophoresis, in athlete rehabilitation, 536 Irrigation, of amputation site, 697

Ischemia, amputation and salvage procedures in, in adults, 667-689 art of, 667-668 internal versus external, 682-684 level for, 668-671, 681-682 postoperative care in, 684-686 procedure for, 678-681 team approach to, 671-672 wound healing considerations in,

668-671 in pediatric patients, 657-661 arterial occlusion, 659 gangrene after clubfoot surgery,

659-660 part strangulation, 660-661 purpura fulminans, 657-658

in diabetes mellitus, 237, 240 magnetic resonance imaging of, 258-259

in peripheral vascular disease, 267-268 Isoflurane, for pediatric patients, 142

Joint space narrowing, in arthritis, radiography of, 371 Jones' fractures, in young athletes, 570

Keller procedure, in hallux limitus, 425 Keratosis. See Hyperkeratosis. Ketamine, for pediatric patients, as anesthetic, 142 as premedication, 137-138

Klippel-Weber-Trenaunay syndrome, macrodactyly in, 644-645, 650 Knee, deformities of, hip instability in, 185 in-toe gait in, 211-212 salvage of, in tibial hemimelia treatment, 633

soft tissue imbalance at, in-toe gait in, 73

Laboratory testing, preoperative. See Preoperative testing. Langer Counter Rotation System splint,

Laryngospasm, perioperative, in pediatric patients, 144

Laser Doppler studies, in healing prediction, 669-670

Lateral compartment syndrome, in peroneal tendon injury, 454 Lawnmower injury, reconstruction of,

followed by amputation, 605-608 Lee procedure, for ankle stabilization, 463, 465

Leriche's syndrome, angiography of, 268-269

Leukocytes, in diabetic foot scan, indium 111-labeled, 249-250 technetium 99m-labeled, 250 Lidocaine, in pediatric surgery, 140

Ligamentous and Capsular Repair System, for arthroscopic ankle stabilization,

Ligamentous laxity, 117-130 causes of, 124-125 clinical features of, 125, 127-128 in adolescents, 123-124 in infants, 120-121 in school-age children, 122-123 in toddlers, 121-122 intrauterine environment and, 118, 120 treatment of, 128 Ligature strangulation, in polydactyly

treatment, 657 Limb lengthening, in tibial hemimelia, 633 Limb salvage, in pediatric patients. See

Pediatric patients, amputation and salvage procedures in. versus amputation, in major trauma,

603-612 Limb Salvage Index, for treatment decisions, 609-610

Lipoma, of calcaneus, imaging of, 293 Lisfranc amputation and modifications, 753-755

prostheses for, 780-781

versus Syme's ankle disarticulation, 763 Lisfranc joint. See Tarsometatarsal joint complex (Lisfranc joint).

Ludloff procedure, in hallux limitus, 424

Macrodactyly, 644-652 peripheral nerve tumors in, 645 simple, 650-652 syndromes associated with, 645, 648-650 treatment of, 650-652 Magnetic resonance imaging, of Achilles

tendinitis and bursitis, 289-291 of arthritis, 395, 397

of calcaneal avascular necrosis, 288-289 of calcaneal fractures, 287-288, 353 of diabetic foot, 252-261

in edema, 253-254 in ischemia, 258-259

in neuroarthropathy, 257-258 in osteomyelitis, 256-257 in soft-tissue fluid collections, 254-256

in surgical changes, 259

of heel pain, avascular necrosis, 288-289 os trigonum syndrome, 297-298 plantar fasciitis, 285-286

rheumatoid arthritis, 295 tumor-related, 293-294

of hip, 193

of Morton's neuroma, 308-309

of tendon injuries. See Tendon injuries. Magnification radiography, of diabetic foot, 243-244

Malformations, congenital, treatment of. See Pediatric patients, amputation and salvage procedures in.

Malignancy, 613-620

amputation in, 615-616 fibrosarcoma, 616

> malignant melanoma, 616-617 margins for, 615

squamous cell carcinoma, 617-619 diagnosis of, 615

initial evaluation of, 614

treatment of, 615-616

Malignant hyperthermia, in pediatric patients, 143-144

Malignant melanoma, treatment of, 616-617

Malleolus, amputation at. See Syme's amputation, transmalleolar. fractures of, in young athletes, 572

Mangled Extremity Severity Index, for treatment decisions, 608-610 Manipulation, in athlete rehabilitation,

538-539 of clubfoot, 32-35

Marfan's syndrome, ligamentous laxity in, 125

Markel supinator shoe, for out-toeing, 212 Massage, in athlete rehabilitation, 538-539 Mau procedure, in hallux limitus, 424 Mayo procedure, in hallux limitus, 425 Medial-lateral flaps, in digital amputation, 709-710

Medial-plantar flaps, in digital amputation, 711, 713

Melanoma, malignant, treatment of, 616-617

Meningococcemia, purpura fulminans in, amputation and salvage procedures in, 657-658

MENS (microcurrent electrical nerve stimulation), in athlete rehabilitation,

Meperidine, in pediatric patients, 138 Metabolic cost, of amputation, 599-602 Metatarsal(s), absence of, in ectrodactyly, 637-642

amputation across. See Transmetatarsal amputation.

fifth, fractures of, in young athletes, 570 first, length of, in hallux limitus, 411 osteotomies of, in hallux limitus, 424-425

in hallux valgus, 439-443 plantar epiphyseal arrest of, in flatfoot, 96

position of, in hallux limitus, 411 shape of, in hallux limitus, 410-411 fractures of, after amputation, 731 lengthening of, in brachymetatarsia,

642-643

short (brachymetatarsia), 642-643 Metatarsal pads, for Morton's neuroma, 495

Metatarsalgia, in young athletes, 566 Metatarsophalangeal joint(s), arthroplasty of, as amputation alternative, 703, 706-707

disarticulation of, 715-717

first, fusion of, in hallux limitus, 425 imaging of, in gout, 379-381 in osteoarthritis, 378 injuries of, in ballet dancers, 529-530

turf toe in, 422, 555-556 limited motion of. See Hallux limitus. neuroarthropathy of, in diabetes melli-

tus, 236 Metatarsus adductus, treatment of, shoes

in. 209-211 Metatarsus primus adductus, treatment of, shoes in, 211

Metatarsus primus elevatus, in hallux limitus, 411

Methohexital, in pediatric surgery, 140-141 Microcurrent electrical nerve stimulation. in athlete rehabilitation, 537

Microdactyly, 652

Midazolam, as premedication, for pediatric patients, 136-137

in pediatric surgery, 140–141 Midfoot amputation, 753–760

indications for, 742-743 level of, 742-744

Midtarsal joints, deformity of, in clubfoot, 20, 23

Mini-Statak Soft-Tissue Attachment Device, for arthroscopic ankle stabilization, 475

Mitek anchoring systems, for ankle stabilization, 471–472, 476

Mixed connective tissue disease, arthritis in, imaging of, 382

Molluscum contagiosum, in athletes, 518-519

Morphine, in pediatric patients, 138 Morton's extension, in hallux limitus, 423

Morton's neuroma, 303–311 clinical features of, 303–304

diagnosis of, 304-305 imaging of, 305-309

in athletes, 489-501

anatomic considerations in, 490–493 clinical features of, 489–490 diagnosis of, 490 etiology of, 493–494

treatment of, 494–500 young, 566, 568

Motor sensory neuropathy, hereditary. See Hereditary motor sensory neuropathy. Muscle relaxants, in pediatric surgery, 144 Mycotic aneurysm, aortic, angiography of,

Mycotic aneurysm, aortic, angiography of, 271–272 Myelomeningocele, hip instability in, 185

Myelomeningocele, hip instability in, 185 Myerson classification, of tarsometatarsal joint complex injuries, 360–361

Nail diseases, in athletes, 516–517 Nanocolloid, technetium 99m-labeled, in diabetic foot scan, 252

Nausea, postoperative, in pediatric patients, 145–146

Navicular, deformity of, in clubfoot, 18 stress fractures of, in young athletes, 570 Necrosis, avascular, of calcaneus, imaging

of, 288–289 of Achilles tendon, in athletes, 451–452 Neisseria meningitidis infections, purpura fulminans in, amputation and salvage

procedures in, 657–658 Neoplasms. See *Tumors*.

Nerve(s), hypertrophy of, in Charcot-Marie-Tooth disease, 101–106 transection of, in amputation, 697 Nerve block in pediatric patients 141

Nerve block, in pediatric patients, 141 Neuroarthropathy, in diabetes mellitus, 235–239

imaging of, 257–258, 383–384 Neurofibroma, of hindfoot, imaging of,

Neurofibromatosis, macrodactyly in, 644–645

Neuroma, Morton's. See *Morton's neuroma*. of hindfoot, imaging of, 293–294

Neuromuscular diseases, hip instability in, 184–185

Neuronal form of hereditary motor sensory neuropathy, 106–108 pathogenesis of, 109–110

Neuropathy, hereditary motor sensory. See Hereditary motor sensory neuropathy.

Nevi, in Klippel-Weber-Trenaunay syndrome, 650

Nilsonne procedure, for ankle stabilization, 463–464

NISSA Score, for treatment decisions, 608–610

Nitrous oxide, for pediatric patients, 142–143

Nonsteroidal anti-inflammatory drugs, in Morton's neuroma, 496 Nothing by mouth guidelines,

preoperative, for pediatric patients, 135

Nuclear medicine scans, of arthritis, 387–389 of calcaneal avascular necrosis, 288

of calcaneal fractures, 287, 353 of diabetic foot, 244, 246–252 gallium 67, 247, 249

indium 111, 249–250 technetium 99m, 246–248, 250–252

of heel pain, calcaneal fractures, 287 plantar fasciitis, 285

of tarsometatarsal joint complex injury, 363–366

Nutrition, in healing prediction, 670

Obstetrical factors, in hip instability, 182 Occlusion, vascular, ischemia in, amputation and salvage procedures in, 657-661

Occupational disorders, heel-spur syndrome, 508

Oligohydramnios, hip instability in, 185 Onion bulb formations, in nerves, in hereditary motor sensory neuropathy, 104

Onychomycosis, in athletes, 516–517 Opioids, in pediatric patients, 138

Orthoses. See also *Shoes*. in amputation, 779–782

digital, 728 in foot deformities, in hereditary motor sensory neuropathy, 113

in hallux limitus, 423

in Morton's neuroma, 495 in ray resection, 728

Ortolani's test, for hip instability, 187–188
Os peroneum syndrome, heel pain in,
296–297

Os trigonum syndrome, heel pain in, 297-298

Osteitis deformans (Paget's disease), heel pain in, 297 Osteoarthritis, imaging of, 377-379 Osteoarthropathy, after amputation, 731, 733-734

Osteochondral fractures, of ankle, in young athletes, 572-573

Osteochondrosis, of calcaneal apophysis (Sever's disease), 298 Osteoclastoma, of calcaneus, imaging of,

Osteogenesis imperfecta, ligamentous laxity in, 125

Osteoid osteoma, of calcaneus, imaging of, 292-293

Osteomyelitis, amputation in, 695 imaging of, 382-384 in diabetes mellitus, 240-241 computed tomography of, 244

imaging of, 383-384 magnetic resonance imaging of, 256-

magnification radiography of, 243-244 nuclear medicine scans of, 244, 246-252

treatment of, 674-677 Osteophytosis, in arthritis, radiography of, 371

Osteoporosis, in arthritis, radiography of, 371

Osteotomy, in amputation, 717 in arthroplasty, 702-703 in ectrodactyly, 639 in flatfoot, of cuneiform, 96 in hallux limitus, in athletes, 424-425 in hallux valgus, in athletes, 438-443 in macrodactyly, 650 Out-toe gait, shoe therapy for, 212-213 Overgrowth, of toes, 644-652

Overuse injuries, in ballet, 525-532 Oxygen measurement, transcutaneous, in healing prediction, 669

PACT (Preservation Amputation Care and Treatment) program, in amputation prevention, 787-790

Paget's disease, heel pain in, 297 Pain, heel. See Heel pain; Heel-spur syndrome.

in ectrodactyly, 640-641 in hallux limitus, 415-416 in Morton's neuroma, 489-490 phantom, in amputation, 589, 591 Painful heel syndrome (plantar fasciitis), 284-287

anatomic considerations in, 283 Palmen's sign, in hip instability, 188 Paraplegia, spastic (hereditary motor sensory neuropathy, type V), 109 Paratendinitis, 314 of Achilles tendon, magnetic resonance imaging of, 317-318, 320 Part strangulation, 660-661 Patella, squinting, in in-toe gait, 76 Pediatric patients, amputation and salvage procedures in, 621-665

in congenital conditions, 621-657 ankle, 621-633 brachymetatarsia, 642-643 classification of, 621-622 constricting band syndrome,

633-638 ectrodactyly, 637-642 fibular hemimelia, 623-624, 626-632 hemihypertrophy, 644-652 macrodactyly, 644-653 microdactyly, 652 polydactyly, 652, 654-657 syndactyly, 652, 654-657 tibial hemimelia, 624-627, 632-633

in limb ischemia, 657-661 arterial occlusion, 659 gangrene after clubfoot surgery, 659-660 part strangulation, 660-661

purpura fulminans, 657-658 analgesia in, 131-148 anesthesia in, 131-148 bracing techniques for, 217-228 Charcot-Marie-Tooth disease in, 99-116 clubfoot in. See Clubfoot. hip instability in, 179-208 in-toe gait in, evaluation of, 57-85 ligamentous laxity in, 117-130 preoperative testing in, 149-178 sedation in, 131-148 shoe therapy for, 209-214 splinting techniques in, 217-228

sports injuries in, 559-579 above ankle, 574-575 ankle, 570-573 apophyseal, 573-574 forefoot, 565-568 healing in, 561 hindfoot, 568-570 midfoot, 568-570 risks for, 563-565 tendon, 573-574 unique anatomy and physiology in, 560-562

subtalar arthrorisis in, 87-98 Peg, polyethylene, in subtalar arthrorisis, 93, 95

Periostitis, in young athletes, 574 Peripheral vascular disease, 265-280 angiography of, 268-273 in adventitial cystic disease, 273

in aneurysmal disease, 270-274

in entrapment, 273

Peripheral vascular disease (Continued) in ergotism, 273 in occlusive disease, 268-270 in thromboangiitis obliterans, 273 in venous disease, 278-279 evaluation of, 266-268 factors affecting, 265-266 management of, 267 prognosis for, 266 risk factors for, 266-267 Peritendinitis, definition of, 314 Perkin's line, in hip radiology, 191-192 Peroneal cuboid syndrome, in athletes, 455 Peroneal muscular atrophy. See Charcot-Marie-Tooth disease. Peroneal tendon, in ankle ligament reconstruction, 463-467 injuries of, in athletes, 453-457 magnetic resonance imaging of, 324tendinitis of, in ballet dancers, 526-527 in young athletes, 574 Pes cavus, heel-spur syndrome with, 506 in Charcot-Marie-Tooth disease, 103 in hereditary motor sensory neuropathy, pathogenesis of, 112 Pes planus. See Flatfoot. Pes valgus, convex, in hip instability,

183–184 shoe therapy for, 212 Petechiae, in athletes, 512–513 calcaneal (black heel syndrome),

511–512 Phalanx, first, osteotomy of, in hallux limitus, 424–425

Phantom sensation and pain, in amputation, 589, 591 Phonophoresis, in athlete rehabilitation,

537–538 Photoplethysmography, in healing

prediction, 669
Physical therapy, for amputation, 591–592,

727–728 in athlete rehabilitation, electrical agents in, 536–537

for ankle sprain, 539–555 for hallux valgus, 443–445 for heel-spur syndrome, 556 for Morton's neuroma, 495–496 for plantar fasciitis, 556 for sesamoiditis, 556 for turf toe, 555–556

for turf toe, 555–556 iontophoresis in, 536–537 mechanical agents in, 537–539 thermal agents in, 534–535

Pigeon toeing. See *In-toe gait*. Piston sign, in hip instability, 189 Plagiocephaly, hip instability with, 184 Plantar fascia, release of, in heel-spur syndrome, 506 Plantar fasciitis, 284–287 anatomic considerations in, 283 in athletes, rehabilitation in, 556 young, 569

Plantar flap, in transmetatarsal amputation, 746–751 Plantar keratosis, in athletes, 522–523

Plaster splints, 227
Plates, gait, for in-toe gait correction, 81–82

Polydactyly, 652, 654-657

Polyethylene peg, in subtalar arthrorisis, 93, 95

Polysyndactyly, 652, 654–657 Popliteal arteries, angiography of, in adventitial cystic disease, 273 in entrapment, 273 in occlusive disease, 270

Porokeratosis plantaris discreta, in athletes, 522

Postaxial polydactyly, 655, 657 Posterior tibial tendon dysfunction, in athletes, 479–488, 574 young, 569

Postoperative care, in amputation, 588–589 complications in, 590 digital, 726–728 for infection or ischemia, 684, 686 Syme's, 766–768

transmetatarsal, 747 in ray resections, 726–728

Post-static dyskinesia, in heel-spur syndrome, 503

Preaxial polydactyly, 654 Predictive Salvage Index, for treatment decisions, 608, 610

Premedications, for pediatric patients, 136–138

Preoperative testing, for amputation, 588 in pediatric patients, **149–178** amount needed, 151 historical review of, 151 medicolegal considerations in, 150 problems with, 150–151 recommendations for, 176 review of 203 cases, 152–175

types of, 149-150 Preservation Amputation Care and Treatment program, in amputation prevention, 787-790

Preventive interventions, for amputation. See *Amputation*.

Prewalker shoes, 209–210 Primary care specialist, podiatrist as, 786

Propofol, in pediatric surgery, 141 Proprioception exercises, in ankle sprain, in athletes, 542–543, 549–552

Prostheses, for amputation, **775–783** ambulation with, biomechanics and pathomechanics of, **775–778**  metabolic cost of, 599–602 Chopart, 780–782 digital, 728, 777–778 for infection, 686 foreshortened shoe, 782 hallux, 777, 779 immediate postoperative fitting of, 590 Lisfranc, 780–781 Syme's, 767–768, 770, 772 transmetatarsal, 778–780 for ray resections, 778 Proteus syndrome, macrodactyly in, 644–645, 648–649

644–645, 648–649
Proximal articular set angle measurement, in hallux valgus, in athletes, 437
Psoriasis, in athletes, 519–521
Psoriatic arthritis, imaging of, 374–376
Psychological issues, in amputation, 595
Pulse evaluation, in healing prediction, 668–669

Puncture wounds, in athletes, 514–515 Purpura, in athletes, 512–513 Purpura fulminans, amputation and salvage procedures in, 657–658

Radiography, of arthritis, 369–385 degenerative (osteoarthritis), 377–379 gouty (crystal deposition), 379–381 in connective tissue diseases, 381–382 neuropathic, 382–384 principles of, 369–370 rheumatoid, 372–374 septic, 382–384 seronegative spondyloarthropathies, 374–377 sesamoid bones, 384

of calcaneal avascular necrosis, 288 of calcaneal fractures, 341–343 of diabetic foot 242–243

of diabetic foot, 242-243 magnification, 243-244

of Haglund's disease, 291–292 of heel pain, avascular necrosis, 288

Reiter's syndrome, 295 of hip instability, 190–192

of Morton's neuroma, 305 of osteomyelitis, 382-384

of Paget's disease, of heel, 297

of tarsometatarsal joint complex injury, 360, 362–363

Radionuclide scans. See Nuclear medicine scans.

Radiotherapy, with amputation, in malignancy, 615

Range of motion exercise, in ankle sprain, in athletes, 539–541 Ray(s), absence of, in fibular hemimelia,

623–624 in tibial hemimelia, 626 excess number of (polysyndactyly), 652, 654–657

resection of, biomechanical effects of,

complications of, 728–735 in macrodactyly, 650 in polydactyly, 655–656 indications for, 695–696

postoperative care in, 726–728 principles of, 696–700

procedures for, 721–726 prostheses for, 778

rehabilitation in, 727–728 tissue stabilization in, 700–701

Reconstruction, versus amputation, in major trauma, 603–612

Recreation, for amputee, 596

Reflex sympathetic dystrophy, heel pain in, 298–299

Refsum's syndrome, 108-109

Rehabilitation, for amputation, ambulation potential and, 594–596 contralateral foot condition in, 593

digital, 727–728 home equipment for, 593–594 physical therapy for, 591–592, 727–728

physical therapy for, 591–592, 727–72 recreational activities in, 596 stump care in, 592–593 support groups in, 597

in athletes, 533-557

after hallux valgus correction, 443–445 electrical agents in, 536–537

exercise in, alternative, 533–534 in ankle sprain, 539–555 in heel-spur syndrome, 556

in plantar fasciitis, 556 in sesamoiditis, 556

in turf toe, 555-556 iontophoresis in, 536

mechanical agents in, 537-539 phases of, 534

thermal agents in, 534–535 Reiter's syndrome, heel pain in, 294–295

imaging of, 374–375, 377
Retinitis pigmentosa, in hereditary motor sensory neuropathy, type VII, 109

Retrocalcaneal bursitis, imaging of, 291–292

magnetic resonance, 319–320 Return to activity exercises, in ankle sprain, in athletes, 543, 551, 554

Reverdin-Green procedure, in hallux valgus, in athletes, 441–443

Rheumatoid arthritis, bone scan of, 389 heel pain in, 295 imaging of, 372–374

RICE treatment, in Achilles tendon injuries, 450

in athlete rehabilitation, 534–535

Ring constrictions, of limbs and digits, 633–636 Rocker-bottom deformity, in diabetes mellitus, 237 Rocker-bottom shoe, after amputation, 780–781 in hallux limitus, 422–423 Roussy-Levy syndrome, 104 Russian stimulation, in athlete rehabilitation, 537

SACH (soft ankle cushion heel) prosthesis, 767–768

Salvage procedures. See Limb salvage; Pediatric patients, amputation and salvage procedures in.

Sand toe, rehabilitation in, 555-556 Sanders classification, of calcaneal

fractures, 351-353 Schizodactyly, 654

Schwannoma, macrodactyly in, 645 Scintigraphy. See Nuclear medicine scans.

Sclerosis, bony, in arthritis, radiography of, 371

Sedation, in pediatric patients, 131–148 as premedications, 136–138 complications of, 143–144 conscious, 132 controversies in, 138–139 deep, 132 definition of, 132 intraoperative, 139–143 nothing by mouth guidelines for, 135 outpatient, 134–135 outside operating room, 132–133

postoperative considerations in, 144–147 preoperative examination for, 133–134

Self-image, in amputation, 587 Sensation, phantom, in amputation, 589, 591

Sensory motor neuropathy, hereditary. See Hereditary motor sensory neuropathy. Sensory neuropathy, in diabetes mellitus, 235–239

Septic arthritis, imaging of, 382–384 Seronegative spondyloarthropathies, imaging of, 374–377

Sesamoid bones, arthritic involvement of, imaging of, 384 injury of, hallux limitus and, 422

Sesamoiditis, in athletes, 566 rehabilitation in, 556

Sever's disease, heel pain in, 298 in young athletes, 573–574 Sevoflurane, for pediatric patients, 142

Shenton's line, in hip radiology, 192 Shin splints, in young athletes, 574 Shock, emotional, in amputation, 589–590 Shoes, 209–214 after amputation, Chopart, 780–782 digital, 728 foreshortened, 782

Lisfranc, 780-781 transmetatarsal and forwa

transmetatarsal and forward, 779-780 ballet, 525, 531

fitting problems with, in ectrodactyly, 637–638

for Achilles tendon injuries, 450 for hallux limitus, corrective, 422–423 examination of, 416–417

for in-toeing, 209-212

for Morton's neuroma, 493-494

for out-toeing, 212–213 for ray resection, 728 prewalker, 209–210

regular last reversed, 210

types of, 213-214

with splinting systems, 223

Simons rule of 15, in clubfoot evaluation, 27

Skin disorders. See *Dermatologic problems*. Skin perfusion pressure measurement, in healing prediction, 669–670

Sleeping position, in ligamentous laxity, 120–121

Smoking cessation programs, in amputation prevention, 792–798 Soft ankle cushion heel prosthesis, 767–768

Soft-tissue, coverage of, in digital amputation, 719–720

fluid collections in, in diabetes mellitus, magnetic resonance imaging of, 254–256

Spina bifida, hip instability in, 185 Splinting and bracing, in pediatric

patients, 217-228 age and, 219-220

correction extent in, 221–222 deformity type and level and, 217–219

Denis-Browne, 1-4, 215-216

in in-toe gait, 77–81 devices for, 223–228 foot stability and, 223

night, 217 shoes for, 223

treatment length in, 220–221 types of, 219

Spondyloarthropathies, seronegative, imaging of, 374–377

Sports medicine, Achilles tendon injuries, 447–458

amputation, 596 ankle arthroscopic stabilization, **459–478** ballet overuse injuries, **525–532** 

dermatologic problems, 511–524 hallux limitus, 407–426

hallux valgus, 427–445 heel pain, 503–509

Morton's neuroma, 489–501

peroneal tendon injuries, 453-458 posterior tibial tendon dysfunction, 479-488

rehabilitation, 533-557

young athletes, 559-579

Sprain, ankle, in athletes, arthroscopic stabilization of. See Ankle, arthroscopic stabilization of. rehabilitation in, 539-555 young, 571-572

Spur, heel. See Heel-spur syndrome. Squamous cell carcinoma, treatment of, 617-619

Stiffness, in hallux limitus, in athletes, 415-416

Stimulation, electrical, in athlete rehabilitation, 536-537

Stone procedure, in hallux limitus, 425 Strangulation, from constricting object, 660-661

Strapping, in ankle sprain, in athletes, 554-555

Streeter's dysplasia (constricting band syndrome), 633-636

Strength exercises, in ankle sprain, in athletes, 541-542, 544-548

Stress fractures, in ballet dancers, 530-531 in young athletes, 566 navicular, 570 tibial, 574

of calcaneus, 287-288

radionuclide scans of, 353 Stump, in amputation, care of, 592-593

Subchondral cysts, in arthritis, radiography of, 371 Subtalar arthrorisis, 87-98 complications of, 97-98

indications for, 88, 90-91 postoperative care in, 97

procedure for, 91-95 Subtalar joint, deformity of, in clubfoot, 20 Succinylcholine, in pediatric surgery, 144 Sudek's atrophy, heel pain in, 298-299 Support groups, for amputee, 597 Surgery, in diabetic foot, magnetic

resonance imaging of, 259 Sutures, for amputation site, 699-700 removal of, 727

Sweating, excessive, in athletes, 515-516 Swelling, in arthritis, radiography of, 371

Syme's amputation, terminal, 707-708 transmalleolar, 763-774 advantages of, 764 anesthesia for, 765-766 complications of, 768-772 history of, 763 in lawnmower injury, 606, 608 indications for, 764-765

> parameters for, 765 postoperative care for, 766-768

two-stage versus one-stage, 764-765 versus other hindfoot amputations, 763-764

Syndactyly, 652, 654-657 Synovial cysts, in athletes, 523 Synovial inflammation, imaging of, 372-377

procedure for, 766

Synovitis, in hallux limitus, 421 Systemic lupus erythematosus, arthritis in, imaging of, 381-382

Talipes equinovarus. See Clubfoot. Talocalcaneal joint, deformity of, in clubfoot, 20

Talocrural joint, deformity of, in clubfoot, 18, 20

Talofibular ligaments, anatomy of, 460 pathomechanics of, 461 reconstruction of, 463-467 repair of, arthroscopic, 468-474 surgical, 467

Talonavicular joint, deformity of, in clubfoot, 20, 23

Talon noir (black heel syndrome), in athletes, 511-512

Talus, deformity of, in clubfoot, 18, 21-22 Tarsal coalition, in young athletes, 569 Tarsal tunnel syndrome, versus heel-spur syndrome, 505

Tarsometatarsal joint complex (Lisfranc joint), injuries of, 357-368 anatomic considerations in, 358-359 biomechanics of, 359 classification of, 360-361 computed tomography of, 367 in diabetes mellitus, 237, 239 late detection of, 364 mechanism of injury in, 359 nuclear medicine scans of, 363-366 overuse, in ballet dancers, 531 radiography of, 360, 362-363 Lisfranc amputation at, 753-755

tus, 236 Taylor bunionette, in young athletes, 568 Team approach, to amputation, 586 to limb salvage, in ischemia or infection, 671-672

neuroarthropathy of, in diabetes melli-

Technetium 99m scans, of arthritis, 388 of diabetic foot, 246-248, 250-252 Telescope sign, in hip instability, 189 Tendinitis, Achilles, imaging of, 289-292 in ballet dancers, 526-528 in young athletes, 574

Tendon(s), disorders of. See specific tendons, e.g., Achilles tendon injuries. transection of, in amputation, 697

Tendon injuries, 313-335 anatomic considerations in, 313-315 magnetic resonance imaging of, Achilles tendon, 317-320 anterior tibial tendon, 329-331 flexor hallucis longus tendon, 328-329 peroneal tendons, 324-328 posterior tibial tendon, 320, 322-324 technique for, 316-317 pathogenesis of, 314 Tendonosis, definition of, 314 of Achilles tendon, magnetic resonance imaging of, 317-319, 321 Tenography, in arthritis, 390, 392-393 Tenosynovial cysts, in athletes, 523 Tenosynovitis, magnetic resonance imaging of, flexor hallucis longus, 328-329 peroneal, 326-327 peroneal, in athletes, 455-456 TENS (transcutaneous electrical nerve stimulation), in athlete rehabilitation, Terminal Syme's amputation, 707-708 Thermotherapy, in athlete rehabilitation, Thiopental, in pediatric surgery, 142 Thromboangiitis obliterans, angiography in. 273 Thrombosis, venous, venography in, 278-279 Tibial arterial disease, angiography of, 270 Tibial fractures, in ballet dancers, 531 in young athletes, 574 Tibial hemimelia, 624-626 treatment of, 626-627, 632-633 Tibial tendon, dysfunction of, in athletes, anterior, 574 posterior, 479-488, 569, 574 injuries of, magnetic resonance imaging of, anterior, 329-331 posterior, 320, 322-324 Tibial torsion, internal, splinting for, 218 in-toe gait in, 68-73 Denis-Browne splints for, 77-80 gait compensations in, 73-75 internal, 68, 70-71 measurement of, 71-73 natural history of, 69-71 normal values of, 69-71 presentation of, 68-69 transverse plane deformities in, 73-77 residual, consequences of, 75

shoe therapy for, 211 Tibial Torsion Transformer, 227

Tibiofibular torsion, 68, 70

Tinea pedis, in athletes, 517-519

TIPC mnemonic, for amputation decisions,

Tibiotalar torsion, 69

693

Tissue stabilization, in amputation, 700-701 Tobacco use, cessation programs for, in amputation prevention, 792-798 Toe(s), absence of, in ectrodactyly, 637-642 amputation of, 691-739 biomechanical effects of, 777-778 complications of, 728-735 definitive, 696 distal, advantages of, 696 incidence of, 691 indications for, 693-696 postoperative care in, 726-728 prevention of, 692-693 principles of, 696-700 procedures for, 701-721 alternatives for, 701-707 dorsal-plantar flaps in, 709-711 for soft-tissue coverage, 719-720 in toe-to-thumb transfer, 720 in trauma, 717-719 medial-lateral flaps in, 709-710 medial-plantar flaps in, 711, 713 metatarsophalangeal joint disarticulation in, 715-717 Syme's, 707-708 trap-door flaps in, 713, 715 prostheses for, 728, 777-778 rehabilitation after, 727-728 tissue stabilization in, 700-701 arthroplasty of, versus amputation, 694-695, 702-707 clawed, in hereditary motor sensory neuropathy, pathogenesis of, 112 constricting band syndrome of, 633-636 crush injury of, reconstruction in, 604denervation of, in macrodactyly, 650 drifting of, after amputation, 735 excess number of (polydactyly), 652, 654-657 hammer, in hereditary motor sensory neuropathy, pathogenesis of, 112 hemihypertrophy of (macrodactyly), 644-652 hypoplastic (microdactyly), 652 infection of. See Infections. ischemia of. See Ischemia. malignant melanoma of, treatment of, 616-617 malposition of, in brachymetatarsia, 642-643 pressure measurement in, in healing prediction, 669, 743-744 sand, rehabilitation in, 555-556 strangulation of, from constricting object, 660-661 turf, hallux limitus and, 422 rehabilitation in, 555-556 webbed (syndactyly), 652, 654-657

Toeing-in. See *In-toe gait*.
Toeing-out, shoe therapy for, 212–213
Toe-to-thumb transfer, amputation for, 720
Tophi, in gout, imaging of, 379–380
Torquheel device, circular, 227–228
Torsion, definition of, 58
Torticollis, hip instability with, 184
Total lymphocyte count, in healing prediction, 670, 765
Tourniquet, for amputation, 746
Transcutaneous electrical nerve

stimulation, in athlete rehabilitation, 537 Transcutaneous oxygen measurement, in

healing prediction, 669, 743, 765 Transmalleolar amputation. See Syme's amputation, transmalleolar.

Transmetatarsal amputation, **741–762** anesthesia for, 745 biomechanical effects of, 778 elective, 742–744 emergent, 742–744 function after, 759–760 indications for, 741–742 level for, 742–744 open (guillotine), 752–753 plantar ulcer excision with, 747, 751–752 postoperative care in, 747

prostheses for, 778 technique for, 746–747 Transplantation, hallux, to hand, 720 Transverse friction massage, in athlete

rehabilitation, 538-539
Trap-door flaps, in digital amputation,

713, 715 Trauma, amputation in, 694

principles of, 746, 748-750

digital, 717–719 major, **603–612** 

case studies in, 604–608 crush injury, 604–605 dilemmas in, 603–604 lawnmower injury, 605–608 predictive guidelines for, 608–610 treatment of, 610–611 Traumeel, in Achilles tendon injuries,

450-451, 453 in peroneal tendon injuries, 456-457

Trendelenburg test and gait, in hip instability, 189, 194–195

Triple hemisection procedure, for Achilles tendon lengthening, 756–759 Tuber angle of Bohler, 339 Tumors, 613–620

amputation in, 615–616 fibrosarcoma, 616 malignant melanoma, 616–617 margins for, 615 squamous cell carcinoma, 617– 619 diagnosis of, 615 imaging of, hindfoot, 292–294 initial evaluation of, 614 treatment of, 615–616 Turf toe, hallux limitus and, 422 rehabilitation in, 555–556 Twister cables, for splinting, 226 Tylomata, in athletes, 522–523

Ulcers, after amputation, 729–731, 776–778
Syme's, 769, 771
amputation in, 776
diabetic. See *Diabetic foot*.
plantar, excision with transmetatarsal amputation, 747, 751–752
Ultrasonography, of hip, 192–193
of Morton's neuroma, 306–308
Ultrasound, in athlete rehabilitation, 537–538
Uni-Bar, 224
Urate crystal deposition diseases, imaging of, 379–381
Urine testing, preoperative, in pediatric

Valenti procedure, in hallux limitus, 425

patients, 171-172

Vascular compromise. See also Peripheral vascular disease. in diabetes mellitus, 237, 240

angiography of, 244–245 magnetic resonance imaging of, 258– 259

Venography, in venous disease, 278–279 Venous disease, venography in, 278–279 Version, definition of, 58, 60 Viral infections, in athletes, 518–519 Vomiting, postoperative, in pediatric patients, 145–146

Wagner formula, for amputation level, in ischemia, 669
Walking. See *Ambulation; Gait*.
Water running, as alternative exercise, in rehabilitation, 533
Waterman-Green procedure, in hallux

limitus, 424

Watson-Jones procedure, for ankle stabilization, 463, 465 Webbed toes (syndactyly), 652, 654–657 Weight distribution, after amputation, 731–732

Weight training, as alternative exercise, in rehabilitation, 533

Wheaton Brace Extender, for in-toe gait correction, 81

Wheaton Bracing System, 226–227 Windswept deformity, splints for, 219

Wound dehiscence, after amputation, Syme's, 768–769

Xanthomatous lesions, in diabetes mellitus, in athletes. 520–521

Y-line (Hilgenreiner's line), in hip radiology, 190

Zeel, in Achilles tendon injuries, 450–451, 453

in peroneal tendon injuries, 456–457 Zwipp classification, of calcaneal fractures, 350–351